

Cytoarchitecture of the Locust Olfactory System


Thesis by

S. Sarah Farivar

In Partial Fulfillment of the Requirements

for the Degree of

Doctor of Philosophy



California Institute of Technology

Pasadena, California

2005

(Defended February 18, 2005)

© 2005

S. Sarah Farivar

All Rights Reserved

Acknowledgements

I would like to thank my thesis advisor, Gilles Laurent, for the opportunity to conduct this research and for his guidance in its progress. I would also like to thank my committee members Erin Schuman, Henry Lester and Christof Koch for advice and encouragement along the way. Thanks to Scott Fraser, his lab and the Biological Imaging Center where I spent many hours doing all of my imaging work.

All of my lab-mates, past and present, have afforded me with delightful scientific conversations and many mostly delightful non-scientific conversations. Thanks to Javier Perez-Orive, Ben Rubin, Stijn Cassenaer, Vivek Jayaraman, and Roni Jortner for allowing me to vent on just about every topic. I'm indebted to Ofer Mazor for being a great office mate, in several offices, and generally introducing me to my husband.

Thanks to all of my fellow biology classmates for a fabulous first-year. The girls of girls'-lunch have my eternal gratitude for their unending support, always showing up for lunch, and our deep lunchtime discussions.

I owe my parents all my thanks for their encouragement and support (as well as much of my nutrition) these past 23 years that I've insisted on staying in school. Thanks to my brother Shervin for his computer expertise and numerous hours spent debugging my laptop. Lastly, thanks to Adam Hayes for his unending criticisms, many late-nights keeping me company in lab, his MATLAB programming expertise and his support of my work and me in general.

Abstract

The insect mushroom body (MB) receives and processes olfactory information. The MB is a highly conserved structure found in all but a few insect species, and has been shown to be a relevant area in learning and memory of olfactory information. The functional properties of the intrinsic cells of the MB — the Kenyon cells (KCs) — have been extensively studied, and their integrative properties are starting to be understood, particularly in locust. To help decipher its role in odor processing, this thesis presents an in-depth study of the architecture of the locust MB, using a variety of anatomical techniques and original software. Four divisions in the MB's input area, the calyx, are defined and described, as well as a division in one of its output regions, the β lobe. KCs are characterized based on their morphologies and extents within the calyx divisions and the β lobe. MB input cells — the projection neurons — are described in relation to their own input area, the antennal lobe, as well as their output to MB calyx divisions. Two classes of cells downstream from the KCs are also defined anatomically and related to immunochemistry on neurotransmitters. A specific area within the brain — the lateral horn lobe — to which projection neurons and extrinsic cells project, is also defined. Similarities of these structures to other insect orders are also discussed.

Abbreviations

PN Projection Neuron

LN Local Neuron

AL Antennal Lobe

MB Mushroom Body

ORN Olfactory Receptor Neuron

OB Olfactory Bulb

KC Kenyon Cell

GABA Gamma-aminobutyric Acid

LH Lateral Horn

LHI Lateral Horn Interneuron

LFP Local Field Potential

EPSP Excitatory Post-synaptic Potential

IPSP Inhibitory Post-synaptic Potential

LY Lucifer Yellow

AF Alexa Fluor

OC Outer-Calyx

MC Mid-Calyx

IC Inner-Calyx

CI Class I Extrinsic Cell

CII Class II Extrinsic Cell

PI Pars Intercerebralis

MOB Main Olfactory Bulb

AOB Accessory Olfactory Bulb

VNO Vomeronasal Organ

Contents

| | |
|--|------------|
| Acknowledgements | iii |
| Abstract | iv |
| 1 Introduction | 1 |
| 1.1 Studying the Sensory Nervous System | 1 |
| 1.2 Olfaction: A General Survey | 2 |
| 1.3 Anatomical Considerations of Olfactory Systems In Insects | 4 |
| 1.3.1 The Antennae and Antennal Lobes | 4 |
| 1.3.2 Mushroom Bodies | 9 |
| 1.4 Examples from Three Insect Orders | 12 |
| 1.4.1 Order Hymenoptera: <i>Apis mellifera</i> | 12 |
| 1.4.2 Order Diptera: <i>Drosophila melanogaster</i> | 15 |
| 1.4.3 Order Dictyoptera: <i>Periplaneta americana</i> | 17 |
| 1.4.4 Comparison Between the Three Orders | 19 |
| 1.5 A Detailed Look at Order Orthoptera: <i>Schistocerca americana</i> | 21 |
| 1.5.1 Olfactory Anatomy: The Antennal Lobes | 22 |
| 1.5.2 Olfactory Anatomy: The Mushroom Bodies | 23 |
| 1.5.3 Olfactory Coding | 26 |
| 1.6 Specific Goals and Layout of this Study | 28 |
| 2 Methods | 29 |
| 2.1 Golgi Staining | 29 |
| 2.2 Intracellular Recording and Dye Fills | 31 |

| | | |
|----------|--|-----------|
| 2.3 | Imaging | 33 |
| 2.4 | Isolating Filled Neurons | 34 |
| 3 | MB Architecture and Innervation | 39 |
| 3.1 | Introduction | 39 |
| 3.2 | Results | 41 |
| 3.2.1 | Calyx Architecture | 41 |
| 3.2.2 | KC morphology | 41 |
| 3.2.3 | PN Axons in the Calyx | 43 |
| 3.2.4 | Peduncular Structure and KC Morphology | 44 |
| 3.2.5 | MB Lobe Structure and KC Morphology | 50 |
| 3.2.6 | MB Structure and Extrinsic Cells | 53 |
| 3.2.7 | Lateral Horn | 56 |
| 3.3 | Discussion | 56 |
| 4 | Brain Atlas | 63 |
| 4.1 | Introduction | 63 |
| 4.2 | Kenyon Cells | 64 |
| 4.2.1 | Kenyon Cells of the Outer-Calyx | 65 |
| 4.2.2 | Kenyon cells of the Mid- and Outer-Calyces | 78 |
| 4.2.3 | Kenyon Cells of the Inner- and Outer-Calyces | 81 |
| 4.2.4 | Dendritic Complementation | 85 |
| 4.2.5 | Innervation of the Pedunculus and Lobes | 88 |
| 4.3 | Extrinsic Cells | 88 |
| 4.3.1 | Class I Extrinsic Cells | 89 |
| 4.3.2 | Class II Extrinsic Cells | 94 |
| 4.4 | Projection Neurons | 101 |
| 4.4.1 | Projection Neurons and the Antennal Lobe | 101 |
| 4.4.2 | Projection Neuron Axonal Output | 118 |
| 4.5 | Discussion | 122 |

| | | |
|----------|---|------------|
| 5 | Concluding Remarks | 124 |
| 5.1 | New Methods of Analysis and Data Presentation | 124 |
| 5.2 | Summary of Results | 125 |
| 5.3 | Discussion | 128 |
| 5.4 | Future Directions | 133 |
| A | <i>nimage</i> Code | 135 |
| | Bibliography | 156 |

List of Figures

| | | |
|-----|---|---|
| 1.1 | (a) Scanning electron micrograph of the tip of an antenna of the locust, <i>S. americana</i> . (b) Higher magnification micrograph of an individual sensillum. Images acquired with the help of Dr. David Barsic. | 5 |
| 1.2 | (a) A Golgi-stained glomerulus of <i>S. americana</i> , showing innervation by a projection neuron (PN) and an olfactory receptor neuron (ORN). For details, see chapter 2. (b) Schematic of ORN convergence from the antenna/nose to the antennal lobe (AL)/olfactory bulb (OB). | 6 |
| 1.3 | Schematic of the connections of the olfactory epithelium (OE) and the vomeronasal organ (VNO) to the main olfactory bulb (MOB) and accessory olfactory bulb (AOB), respectively. Glomeruli in the AOB are smaller; mitral and tufted cells of the AOB project dendrites to several glomeruli in contrast to the one glomerulus innervated per mitral/tufted cell in the MOB. Adapted from [75]. | 7 |
| 1.4 | Schematic of the basic circuitry of the mammalian MOB. Periglomerular cells (PC), short axon cells (SA) and granule cells (GC) are all inhibitory cells. M; mitral cell, T; tufted cell (two types are shown, both marked by T). There are one-way and reciprocal synapses between branches of M and T cells with inhibitory neurons. Adapted from [80]. | 8 |

- 1.5 Schematic of the locust brain, showing the positions of the AL, mushroom body (MB) neuropils (the β lobe [β], the α lobe [α], pedunculus [P], and calyx), Kenyon cell somata (KC), the antennocerebral tract (ACT), the lateral protocerebrum (LP), and the pars intercerebralis (PI). Left hand corner shows the axes of the brain. A: anterior; Po: posterior; V: ventral; D: dorsal. Adapted from [8]. 9
- 1.6 Reconstruction of a PN (see chapter 2) superimposed on right hemisphere of brain schematic from figure 1.5; scale bar is 200 μm . Left hemisphere shows a three-dimensional reconstruction of the MB and a KC (green) within it. Inset (adapted from [28]) shows a drawing of a typical local neuron in the AL. Scale bar is 100 μm 10
- 1.7 Schematic of the brain of the honeybee, *Apis mellifera*. Brain nomenclature is shown on the right hemisphere of the brain. The three divisions of each of the two calyces per hemisphere, the lip (Li), the collar (Co), and the basal ring (BR), feed into the pedunculus (P) and down into the β lobe (β), and the α lobe (α) which projects out of plane of the page. Representation of the calyx zones within the α lobe are schematized in the left hemisphere. The Li, Br, and Co feed to the colored anterior portions of the α lobe, while clawed KCs from all the calycal regions feed to the posterior γ division of the α lobe. s l pr: superior lateral protocerebrum; i l pr: inferior lateral protocerebrum. Adapted from [87]. 14

- 1.8 Schematics of the brain of the fruitfly, *Drosophila melanogaster*. **(a)** The entire brain is depicted, showing the calyces and MB lobe divisions in each hemisphere of the brain. PN axons project through two tracts: the inner antennocerebral tract (iACT) and the medial antennocerebral tract (mACT) to the calyx and the lateral horn (LH). Adapted from [35] **(b)** Representation of the calyces by the KCs within the different subdivisions of the lobes based on morphology and molecular epitope. The calyx is viewed from the dorsal surface of the brain in each of the five instances. The KCs projecting to the γ lobe innervate all parts of the calyx. Adapted from [89]. 16
- 1.9 Schematic of the brain of the cockroach, *Periplaneta americana*. AL PNs project to the calyces via the iACT and terminate in the LH. On the left hemisphere, a sagittal section of the β lobe is shown in colors that are described in the figure 1.10. op: optic lobe; l ca: lateral calyx; m ca: medial calyx. Adapted from [58]. 19
- 1.10 Schematics showing the calycal divisions within the cockroach brain and their representation within the β lobe. **(a)** On the left is a frontal (viewed from ventral surface) representation of the two calyces of one MB showing their various divisions. Each calyx (l ca and m ca) is further subdivided into the outer hemicalyx (o h ca) and the inner hemicalyx (i h ca). Within each hemicalyx is shown the three afferent divisions of the calyx, zones I-III A. On the right, the same divisions are viewed from the top (anterior surface) of the brain, stretched along its anteroposterior axis (A: anterior; P: posterior). Adapted from [93]. **(b)** A sagittal cross-section of the β lobe as described in figure 1.9, showing the representation of the calyces as defined by immunochemistry and Golgi stains. The lateral and medial calyces are represented on the anterior and posterior portions of the lobe, respectively. Further, the zones are represented in order from one side to the other. The γ lobe contains clawed KCs from all calycal divisions. Adapted from [94]. 20

| | | |
|-----|---|----|
| 2.1 | A flowchart of the <i>nimage</i> algorithm. The cube outlined in black represents the image stack. p in step 1 is the starting pixel. In step 2, the yellow shell around p determines pixels to consider as background for comparison to p . In step 3, the blue cube B defines the neighbors of p to be searched. In step 4, two pixels in B were searched and are found ‘hot’ using 2. Step 5 shows the neighbors of those two pixels that will be searched next. The algorithm continues to search neighbors of hot pixels until no more hot pixels are found and no more p are entered by user. | 36 |
| 2.2 | Ray-traced reconstructions of a pair of KCs filled with LY. (a) Ventral and lateral views of the reconstructed raw data. (b) Ventral and lateral views of the reconstructed <i>nimage</i> extracted data. | 38 |
| 3.1 | Three-dimensional Imaris reconstruction of the locust calyx showing newly defined divisions, depicted in three views. From left to right, facing out of page, are the ventral, anterior and lateral views (or front, top and side views respectively). The outer-calyx (OC) is depicted in yellow, the mid-calyx (MC) in orange, the inner-calyx (IC) in green, and the accessory calyx in purple. The dorsoventral axis is depicted by ‘D’ at the dorsal end and ‘V’ at the ventral end. The mediolateral axis is depicted by ‘M’ and ‘L’. The anteroposterior axis is depicted by ‘A’ and ‘P’. Scale bar is equal to 70 μm | 42 |

- 3.2 Ray-traced reconstructions of two non-OC-exclusive KCs viewed from the ventral surface of the brain. These same two cells are reconstructed in the following figures within their respective calyx structures. **(a)** An OC/MC cell. Small cluster of MC stubby dendrites are indicated by the red arrow. Scale bar is equal to 15 μm . **(b)** An OC/IC cell. Cluster of IC projections is indicated by the red arrow. Note that the cell body is not shown; the green star indicates the primary process leading to the soma; the green arrow indicates the KC axon. Scale bar is equal to 30 μm 43
- 3.3 Three-dimensional Imaris reconstruction of a KC with branches in the IC (arrow) and OC. All calyx neuropil divisions are rendered in the same color. The same three views are depicted as in figure 3.1 - from left to right: front, top and side views. Scale bar is 100 μm 44
- 3.4 Three-dimensional Imaris reconstruction of half of the locust calyx showing a KC that innervates the mid-calyx and the outer-calyx. Left image shows the frontal view of the calyx with three divisions, in different colors. Black box shows the area of the calyx reconstructed in the right two images. Middle image shows the KC reconstructed with all calycal divisions: OC in yellow, MC in orange and IC in green, from the front view. The green asterisk shows the short MC dendrites just outside the OC. Right image shows the KC reconstructed with the MC only, viewed from the top, with the green asterisk again showing the position of the MC dendrites. 45
- 3.5 Golgi-stained frontal sections of the locust MB depicting calycal divisions. **(a)** Lateral side of the calyx with a red star indicating the position of the IC. A clear division can be seen between this and the outer-calyx. **(b)** A superficial section of the calyx where the thin MC can be seen. Part of the MC is outlined by the red rectangle. 45

- 3.6 Imaris reconstruction of the calyx and two PNs innervating it, viewed from the anterior or bottom aspect. On the left, the OC is shown in yellow, the MC in orange, the IC in green and the accessory calyx in purple. The right image shows PNs in red innervating the OC exclusively. . . . 46
- 3.7 Calyx division relationship to the pedunculus. Each row shows the calyx and posterior portion of the pedunculus from the ventral or back aspect(left column) and posterior or top(right column) aspect. Top row shows the regions of the calyx (OC in yellow and MC in orange) that feed into the two peripheral fiber bundles (also in orange). The middle row shows the area of the central bundle that the IC feeds into, and the bottom row shows the accessory calyx (purple) and its peduncular innervation. 47
- 3.8 Imaris reconstruction of the pedunculus and β lobes viewed from the dorsal aspect. Three black arrows point to the three fiber bundles that emerge from the calyx. Green arrow shows the point at which the fibers join together. 48
- 3.9 Slightly oblique 15 μm Golgi-stained section, showing a comparison of KC axons in the periphery of the pedunculus vs. the core. (a) Red arrow indicates peripheral spiny axons and green arrow indicates the core smooth axons. Scale bar is 20 μm . (b) High magnification images of the spiny peripheral axons (top) and smooth core axons (bottom). Scale bar is 5 μm 49
- 3.10 Three-dimensional reconstructions of the β lobe. Left image shows the two divisions of the β lobe based on innervation by spiny axons (green division) and smooth axons (red division). Right image shows the processes of a class II extrinsic cell, preferentially innervating only the spiny axon zone, outlined with dotted line. Scale bar is equal to 50 μm . . . 50

| | | |
|------|---|----|
| 3.11 | KC axon morphology within the β lobe. (a) Imaris reconstruction of the β lobe viewed from the black arrow shown figure 3.10. White dotted line shows the position of the Golgi section in (b). (b) Golgi-stained section of the β lobe showing spiny axons at the medial edge and smooth axons at the posterior edge. | 51 |
| 3.12 | The spiny axon division of the β lobe exhibits a non-fluorescent core which KCs do not innervate. Scale bars are 50 μm . (a) 15 μm Golgi section showing a superficial slice of the spiny axon division. Red arrow indicates core. (b) 3 μm optical slice of intrinsic fluorescence showing the beta lobe at the same angle as in part (a). Red arrow indicates core. | 51 |
| 3.13 | Imaris reconstruction of two CII extrinsic cells within the beta lobe. Black arrow indicates the core of the spiny axon division shown in figure 3.12. P: peduncular processes; α : α lobe processes. | 52 |
| 3.14 | Schematic of the locust brain, showing the position of a class I extrinsic cell. Ray-traced reconstruction of a single LY-filled cell overlaid on a schematic of the locust brain. Soma lies in the pars intercerebralis, between the two MB calyces. The cell's processes innervate the MB lobes and the posterior end of the pedunculus. An extension of the β lobe process innervates the LH. Sparse fibers are seen in the midbrain area. Scale bar is equal to 200 μm | 54 |
| 3.15 | Schematic of the locust brain, showing the position of a class II extrinsic cell. The cell image is a ray-traced reconstruction of one cell filled with the dye LY, overlaid on a brain schematic. Soma lies in the LH, and processes are sent to the MB lobes, pedunculus, IC and MC. Diffuse processes are also seen in the LH, off the cell's primary neurite as it passes to the MB. Scale bar is equal to 200 μm | 55 |
| 3.16 | Reconstruction of a pair of CII cells (green) along with the MB neuropil (in red), from the frontal view. Extensive innervation of both lobes and pedunculus is visible. LH, MC and IC innervation are also visible. Scale bar is equal to 100 μm | 59 |

| | | |
|------|---|----|
| 3.17 | 1 μm confocal image of the calyx from a stack of the pair of the LY-filled CII cells depicted in figure 3.16. Red arrows indicate the processes of the CII cells innervating the IC. Scale bar is equal to 100 μm | 60 |
| 3.18 | 20 μm thick ray-traced projection of several images from a confocal stack of the β lobe of the pair of filled CII cells depicted in figure 3.16. Pairs of red arrows indicate pairs of identical fibers from each cell. Scale bar is equal to 25 μm | 60 |
| 3.19 | Ray-traced reconstruction of three PNs filled with LY, showing the entirety of processes with innervated neuropils labeled. Frontal view. . . | 61 |
| 3.20 | Three-dimensional Imaris reconstruction of the set of PNs in 3.19, including the OC and the LH. The red bulb that is labeled as the LH is a defined area visible autofluorescently, and in both Bodian and Golgi stains. This same bulb is innervated by both PN terminals (as shown) and extrinsic cell processes. | 62 |
| 4.1 | Comparison of the three types of OC-exclusive cells, shown from the angle that best depicts classification. (a) An OC-exclusive fan-shaped cell viewed from the front or ventral aspect. This cell is also shown in figure 4.2. (b) An OC-exclusive bipartite cell viewed from the side or lateral aspect. This cell is also shown in figures 4.5(b) and 4.6(b). (c) An OC-exclusive single-tree cell viewed from the front or ventral aspect. This cell is also shown in figure 4.12(b). | 66 |
| 4.2 | (a) Imaris reconstructions of an OC-exclusive, fan-shaped cell from the front, top and side views, from left to right, respectively. (b) Ray-traced reconstruction of the cell in part (a) in the same three views. Green stars indicate the primary neurite from the soma (soma not shown) and green arrows indicate the axon. | 67 |

| | | |
|------|---|----|
| 4.3 | Ray-traced reconstructions of OC-exclusive fan-shaped cells in three views (as before). (a) Green arrows indicate axon. Note the cluster of somata indicated by green stars. (b) Green stars indicate primary neurite from soma (soma not shown) and green arrows indicate the cell's axon. | 68 |
| 4.4 | Ray-traced reconstructions of OC-exclusive fan-shaped cells in three views (as before). (a) Green stars indicated primary neurite from soma (soma not shown) and green arrows indicate the cell's axon. (b) Green stars indicate the soma and green arrows indicate the cell's axon. . . . | 69 |
| 4.5 | (a-c) Imaris reconstructions of OC-exclusive bipartite cells from front, top and side views (left to right). Ray-tracings of these three cells are shown in the following figure. | 70 |
| 4.6 | (a-c) Ray-traced reconstructions of the OC-exclusive cells depicted in figure 4.5 in same three views as before. Green stars indicate the primary neurite from the soma (somata not shown), and green arrows indicate the axon. | 71 |
| 4.7 | Ray-traced projections of an OC-exclusive bipartite cell, from three views. Green stars indicate the soma (a cluster of somata were stained, though only one cell's processes are visible). Green arrows indicate axonal projections. | 72 |
| 4.8 | (a-c) Ray-traced reconstructions of OC-exclusive bipartite cells in three views. Green stars indicate the primary neurite to soma (somata not shown) and green arrows indicate a cell's axonal projection. | 73 |
| 4.9 | (a-d) Ray-traced reconstructions of OC-exclusive bipartite cells in three views. Green stars indicate the primary neurite to soma (soma is shown for only the cell in figure (d)) and green arrows indicate a cell's axonal projection. | 74 |
| 4.10 | (a) Imaris reconstruction of an OC-exclusive single-tree cell in two views. (b) Ray-traced projection of a single-tree cell. | 75 |

| | | |
|------|--|----|
| 4.11 | Ray-traced reconstructions of OC-exclusive single-tree cells in three views. Green stars indicate the primary neurite to soma (somata not shown) and green arrows indicate a cell's axonal projection. | 76 |
| 4.12 | (a-c) Ray-traced reconstructions of OC-exclusive single-tree cells in three views. Green stars indicate the primary neurite to soma (somata not shown) and green arrows indicate a cell's axonal projection. | 77 |
| 4.13 | The mid-calyx in relation to the outer-calyx (yellow). The MC (orange) is shown connected to the parts of the pedunculus to which it projects. Left hand figure is viewed from the dorsal or back aspect; right hand figure is viewed from the posterior or top aspect. Note that the IC and accessory calyx are not depicted. | 78 |
| 4.14 | (a) Imaris reconstructions of an OC/MC cell, reconstructed with outer-calyx (yellow), mid-calyx (orange) and inner-calyx (green). Green star indicates MC processes of the cell. (b) Ray-traced projection of the cell in part (a) in three views. Red arrow indicates mid-calyx processes, and green arrows indicate the cell's axon. | 79 |
| 4.15 | Ray-traced projection of three OC/MC cells. Green stars indicate the primary neurite from the soma (somata not shown), green arrows indicate the cell's axon and red arrows indicates MC projections/varicosities. | 80 |
| 4.16 | Imaris reconstruction showing the divisions of the calyx; IC is green, MC is orange and OC is yellow. Left hand figure is a view from the dorsal or back aspect of the brain and right hand figure is a view from the posterior or top aspect. | 81 |
| 4.17 | (a) Imaris reconstructions of an OC/IC cell in three views. Black arrows indicate the cell's IC processes. (b) Ray-traced projection of the cell depicted in (a). Green stars indicate primary neurite from the soma, green arrows indicate the axon, and red arrows indicate the cell's IC processes. | 82 |
| 4.18 | (a) Imaris reconstruction of an OC/IC cell in three views. | 83 |

| | | |
|------|--|----|
| 4.19 | Ray-traced projection of another OC/IC cell in three views. Green stars indicate the primary neurite (soma not shown), green arrows indicate the axon, and red arrows indicate the cell's IC processes. | 84 |
| 4.20 | Imaris projection of three KCs stained concurrently in one brain, from the front, top and side views. All three cells depicted here are OC-exclusive cells. | 85 |
| 4.21 | Imaris reconstruction of three cells filled in one brain. Green cell (left-most cell in front and top views) is an OC/IC cell, while the other two cells are OC-exclusive cells. | 86 |
| 4.22 | Imaris reconstruction of two KCs filled in one brain. Red cell is an OC/IC cell, and green cell is an OC-exclusive bipartite cell. | 86 |
| 4.23 | (a-b) Imaris reconstructions of two pairs of OC-exclusive cells. Due to their close resemblance, the two cells in each pair were not separable. (c) Ray-traced reconstruction of the side view of the cells shown in part (a). Red arrows indicate the two different cells. | 87 |
| 4.24 | A ray-traced projection of a class I extrinsic cell. | 90 |
| 4.25 | Ray-traced projection of the heel region of the MB, showing six branches (red arrows) of a CI extrinsic cell, corresponding to the NADPH diaphorase positive zones shown in figure 4.27. | 91 |
| 4.26 | Ray-traced projection of the LH; CI terminals in the LH lobe are orange and indicated by the red arrow. | 91 |
| 4.27 | NADPH diaphorase staining in the brain of the locust. Red arrows indicate two of six immunoreactive lamina that begin in the peduncular heel and continue up into the α lobe. Adapted from [67]. | 92 |

| | | |
|------|--|----|
| 4.28 | Odor responses of a class I extrinsic cell. (a) Ten trials with one odor (octanol). Each line (in blue) is one 10 second trial, where the odor was presented from between seconds 2 and 3 (as indicated by red line). Horizontal scale bar is equal to 1 second, vertical scale bar is equal to 20 mV. (b) Histograms of spike counts over ten trials with each of four odors: hexanol, mint, octanol and cherry. Each bin is the sum of 200 mS bins, odors presented during bins 5-10. Horizontal scale bar is equal to 1 second, vertical scale bar is equal to 10 spikes. | 93 |
| 4.29 | CII extrinsic cells. (a) Ray-traced projection of a single CII extrinsic cell, showing its soma in the lateral horn (LH) and its innervation of the MB lobes and pedunculus. (b) Ray-traced projection of a pair of CII cells filled concurrently. | 96 |
| 4.30 | Imaris reconstruction of the pair of CII extrinsic cells shown in figure 4.29(b). The MB neuropil (α and β lobes and calyx) is reconstructed in red. Note that this is a ventral view, but the brain is tilted back slightly. | 97 |
| 4.31 | 6 μ m thick ray-traced projection of the β lobe processes of the pair of CII extrinsic cells shown in figures 4.29(b) and 4.30. Red arrows indicate the processes that branch off at regular intervals into the lobe. | 98 |
| 4.32 | Peduncular and calycal innervation of CII extrinsic cells. (a) Red arrow indicates the fine fibers innervating the pedunculus. (b) Peduncular and calyx innervation of the same cells as in part (a). Red arrows indicated the MC and IC divisions. | 98 |
| 4.33 | Comparison of CII extrinsic cell fill with Bodian stained processes. (a) A 10 μ m Bodian slice showing a bundle of processes entering the MB lobes that correspond to the CII cell processes. (b) A high magnification 10 μ m ray-projection of the primary neurite of the CII cell innervating the MB lobes. | 99 |

| | | |
|------|--|-----|
| 4.34 | Comparison of CII extrinsic cell fill with Bodian stained somata. (a) A 10 μm Bodian slice showing possible candidates for CII somata in the lateral horn. (b) A 10 μm ray-projection exhibiting the soma of a CII extrinsic cell in the lateral horn. | 99 |
| 4.35 | Ray-traced projection of a brain stained for GABA immunoreactivity. Red arrow indicates a cluster of GABA immunoreactive cells that correspond in position to CII somata position. | 100 |
| 4.36 | Imaris reconstruction of the antennal lobe (yellow) showing the position of the projection neuron somata (red). Left image shows the frontal view, with the position of the antennal nerve indicated by the black arrow. Right image shows a side view in which the antennal nerve is in back. The black star indicates the position of the antennocerebral tract. | 102 |
| 4.37 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. Green stars indicate the cell's soma and green arrows indicate the axon. | 103 |
| 4.38 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. Note that there are two somata filled, but only one cell's processes are visible. (b) Ray-traced projection of the cell in part (a), from the same two views. | 104 |
| 4.39 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. Note that the soma is missing. | 105 |
| 4.40 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. | 106 |
| 4.41 | Ray-traced projections of two different PNs from the same two views as before. Note that both cells are missing their somata. | 107 |

| | | |
|------|---|-----|
| 4.42 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. Note that there are two somata filled, but only one cell's processes are visible. | 108 |
| 4.43 | (a) Imaris reconstruction of a single PN within the AL from a slightly oblique frontal view (left image) and the posterior (right image) view. (b) Ray-traced projection of the cell in part (a), from the same two views. Note that the soma is missing. | 109 |
| 4.44 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. Note that the soma is missing. | 110 |
| 4.45 | (a) Imaris reconstruction of a single PN within the AL from the frontal (left image) and posterior (right image) views. (b) Ray-traced projection of the cell in part (a), from the same two views. Green arrows indicate the axon, and the green stars indicate the soma. Note that this PN's soma is in a different location than the other PNs shown above. | 111 |
| 4.46 | Ray-traced projections of two different PNs from the same two views as before. Note that the cell in part (b) is missing its soma. | 112 |
| 4.47 | A pair of PNs filled concurrently, showing different radii in the same AL. These are the PNs shown in figures 4.38 and 4.42. (a) The same pair shown in an Imaris reconstruction from the frontal and posterior views. (b) Ray-traced projections of the cells shown in (a), in the same two views. Overlaps or areas where cells could not be separated are shown in yellow. | 113 |
| 4.48 | A pair of PNs filled concurrently, showing different radii in the same AL. Green PN is also shown in figure 4.37. (a) The pair of PNs shown in an Imaris reconstruction, from the frontal and posterior views. (b) Ray-traced projections of the cells shown in (a), in the same two views, but note that the posterior image is slightly turned about a vertical axis. | 114 |

| | | |
|------|---|-----|
| 4.49 | A pair of PNs filled concurrently, showing different radii in the same AL. These are the PNs shown in figures 4.39 and 4.44. (a) The same pair shown in an Imaris reconstruction from the frontal and posterior views. (b) Ray-traced projections of the cells shown in (a), in the same two views. Overlaps or areas where cells could not be separated are shown in yellow. Note that cell bodies are missing. | 115 |
| 4.50 | (a) Ray-traced projection of three PNs in one AL. (b) Ray-traced projection of three PNs in one AL. (c) Ray-traced projection of two PNs in one AL. | 116 |
| 4.51 | Three cells with the same radius size. (a) Ray-projection of three PNs stained concurrently within one AL. Two are shown in green and one is shown in red. Scale bar is equal to 70 μm . (b) Close up of the area around one set of glomeruli for cells in part (a). The two green cells are visible as different cells because one is less bright than the other (indicated by white arrows). The two cells innervate the same glomerulus. Scale bar is equal to 35 μm | 117 |
| 4.52 | Calyx innervation of PN axons. Imaris reconstructions of PNs within the OC, viewed from the ventral and posterior aspects. (a) Single PN in the OC. (b) Two PNs in the OC. (c) Three PNs in the OC. | 119 |
| 4.53 | Relationship of PN axons and KC dendrites within the calyx. Left hand figure shows an Imaris reconstruction of both cells within the calyx. Right hand figure shows a high magnification ray-projected image of the same set of cells; KC is shown in green, PN is shown in red, and areas of overlap appear in yellow. | 120 |
| 4.54 | Relationship of the lateral horn lobe to the calyx. (a) Lateral view of Imaris reconstructed of a PN (as shown in figure 4.52[c]) within the calyx and LH lobe. (b) Dorsal view of the same reconstruction. | 120 |
| 4.55 | 10 μm thick Bodian-stained section showing the bundle of PN fibers (red arrow) as they innervate the LH lobe (red star). | 121 |

| | | |
|-----|--|-----|
| 5.1 | Schematic of the locust antennal lobe showing innervation pattern in one plane. Dark blue and green cells innervate the same glomeruli and lie within the same plane. Light blue cell lies in the plane with the other two cells, but innervates glomeruli at a different radius. | 126 |
| 5.2 | Schematics of the proposed structure of the MB and its KC innervation. (a) Frontal view of the calyx, showing the pattern of innervation of an OC-exclusive cell (red), an OC/MC cell (blue) and an OC/IC cell (green). It is possible that OC-exclusive cells innervate the spiny axon division of the β lobe (shown in gray). (b) Two more views of the MB; top schematic is the calyx viewed from above (topmost arrow in part [a]), showing representative innervation of the three types of KCs. Bottom schematic is a view of the β lobe (bottom arrow in part[a]) showing the spiny axon and smooth axon divisions within the lobe. It is possible that OC/IC cells innervate the smooth axon division of the β lobe. Note also the core of the spiny axon division, as described in chapter 3. | 127 |
| 5.3 | Schematics of the proposed structure of the MB and its class II extrinsic cell innervation. (a) Frontal view of the MB, with the areas of CII innervation shown in yellow. (b) Two more views of the MB; top schematic is the calyx viewed from above (topmost arrow in part [a]), showing representative innervation of the CII extrinsic cells. Bottom schematic is a view of the β lobe (bottom arrow in part[a]), showing the spiny axon and smooth axon divisions within the lobe. CII extrinsic cells innervate the lobe in a characteristic 'C' shape around the spiny axon division's core, in slightly offset areas. Representative of one cell is shown in yellow while another is shown in orange. There are postulated to be 9 to 12 CII cells covering the spiny axon division. | 128 |
| 5.4 | Schematics of the MB showing the proposed structure of the peduncular fibers. | 131 |

| | | |
|-----|---|-----|
| 5.5 | Frontal schematic of the MB showing the region (in gray) of the α lobe that contains the axon collaterals of KCs which innervate the smooth axon division of the β lobe. | 132 |
|-----|---|-----|

List of Tables

| | | |
|-----|--|----|
| 1.1 | Comparison of olfactory system neuron numbers per half brain | 13 |
| 2.1 | Golgi Solutions | 30 |
| 2.2 | Physiological Saline for Locust | 32 |
| 3.1 | Characteristics of Extrinsic Cell Classes | 53 |